

Interpretation of the Underground Muon Charge Ratio

Draft 1.3

M. C. Goodman, J. Reichenbacher, and P. A. Schreiner

December 12, 2008

Abstract

The MINOS experiment has observed a rise in the underground muon charge ratio $r_\mu = \mu^+/\mu^-$. This ratio can be related to the atmospheric production ratios of π^+/π^- and K^+/K^- . Our analysis indicates that the relevant variable for studying the charge ratio is $E_\mu^{surface} \cos \theta_z$, rather than $E_\mu^{surface}$. We compare a simple energy dependent parameterization of the rise in the charge ratio with more detailed previously published Monte Carlo calculations. We also discuss the importance of two previously neglected effects in this context: the charge sign dependency of the dE/dx in rock, and the energy dependence of heavy primaries on the K^+/K^- ratio.

1 Introduction

1.1 Importance of charge ratio measurements

Atmospheric muons come from the decay of π s and K s produced in hadronic showers when cosmic rays interact in earth's atmosphere. These muons have been studied with energies ranging from hundreds of MeV to well over a TeV. A quantitative understanding of cosmic ray muons has value for a number of diverse topics, from atmospheric neutrinos to the chemical composition of the highest energy cosmic rays. The charge ratio of cosmic ray muons has been previously measured over three orders of magnitude in energy. Recently the MINOS experiment[1] presented data that for the first time showed a rise in the measured charge ratio

$$r_\mu \equiv \frac{N^{\mu+}}{N^{\mu-}} \quad (1)$$

from previous measurements at high values of E_μ or more specifically, high values of $E_\mu^{surface} \cos \theta$. In this paper, we discuss some of the issues involved in the measurement and interpretation of the muon charge ratio. In particular, we develop a simplified model where the rise in the charge ratio can be understood from the properties of π and K mesons, and the observation of the rise can be used to determine the π^+/π^- and K^+/K^- ratios. We also address several other issues related to the charge ratio, including the role of muon energy loss, a detector's Maximal Detectable Momentum (MDM) and the effect of possible differences in the spectral index of cosmic ray Hydrogen and Helium on the interpretation.

Cosmic ray muons in the atmosphere come from the decays of cosmic ray π s and Ks. Since the primary cosmic rays are mostly positively charged protons, more π^+ are expected than π^- . The quark content of the protons and atmosphere has been used to estimate the π^+/π^- ratio to be near 1.27 [2]. The charge ratio for Ks is even higher due to the phenomenon of associated production. Strange particle production starts with the creation of an s quark and an \bar{s} quark. An s quark which ends up in a nucleus is associated with an \bar{s} quark in a $K^+(\bar{s}u)$. The \bar{s} quark will not be in a baryon. There is also K^+K^- pair production. Phase space favors hadronic production of $K^+\Lambda$ over K^+K^- pairs at all energies, so large K^+/K^- ratios are expected.

A standard parametrization of the atmospheric muon energy spectrum is given by Gaisser[3]:

$$\frac{dN_\mu}{dE_\mu} = \frac{0.14E_\mu^{-2.7}}{\text{cm}^2 \text{ s sr GeV}} \times \left\{ \frac{1}{1 + \frac{1.1E_\mu \cos \theta}{\epsilon_\pi}} + \frac{\eta}{1 + \frac{1.1E_\mu \cos \theta}{\epsilon_K}} \right\} \quad (2)$$

The two terms inside the curly bracket give the contributions of charged pions and Kaons. For the hadron mass (m_i) and lifetime (τ_i) and the atmospheric scale height ($h \sim 6.4$ km), the values of $\epsilon_i = m_i ch/\tau_i$ are the energies where the probability of meson interaction in the atmosphere and decay are equal: $\epsilon_\pi=115$ GeV and $\epsilon_K= 850$ GeV[4] The zenith angle is denoted by θ . In those experiments with a flat overburden, the largest muon intensity is at $\cos(\theta) = 1$. Reference [3] gives $\eta = 0.054$, which normalizes the relative importance to the muon flux from π and K decay, and η depends upon the π/K ratio, branching ratios, and kinematic factors which arise due to differences between the π and K masses.

The ratio between the two terms in Equation 2 quantifies the relative contribution to the muon flux from muons due to π and K decay. This ratio depends on the variable $E_\mu^{surface} \cos \theta$ and rises from $\eta = 5.4\%$ at low energies to $\eta \times \epsilon_K/\epsilon_\pi = 40\%$ at high $E_\mu^{surface} \cos \theta$. This substantial rise is caused by the difference in the decay kinematics of pion and Kaon decay, not to any increased amount of Kaon production from primary cosmic ray interactions in the atmosphere.

MINOS provides the first high statistics measurements of the muon charge ratio at $E_\mu^{surface} \cos \theta > 115$ GeV. We will show that this gives the needed sensitivity for extracting information about cosmic ray produced π s and Ks separately. We also note that a similar calculation for neutrinos shows that Ks are the dominant parent for TeV neutrinos, which are the largest backgrounds for astrophysical source searches at neutrino telescopes.

The organization of this paper is as follows: In the following section, we review some previous measurements of the atmospheric muon charge ratio. In Section 3, we discuss some of the issues involved in measuring the charge ratio in a magnetic detector underground, using MINOS as an example. In the next section, we address in particular issues related to the muon energy loss and how the muon energy at an underground detector is related to the energy in the atmosphere. In Section 5, we use Equation 2, which was based on ideas developed by Zatsepin[4], to parameterize the charge ratio in a simplified model. In particular, Zatsepin's work is the earliest reference that we have found which defines the roles of the critical energies $\epsilon_\pi = 115$ GeV and $\epsilon_K = 850$ GeV. There are two important implications from this model. First, the charge ratio depends not simply on the muon energy $E_\mu^{surface}$ but on the combination of the energy and zenith angle $E_\mu^{surface} \cos \theta$. Second, together with previous measurements at low $E_\mu^{surface} \cos \theta$, observation of the rise in the charge ratio, which occurs between ϵ_π and

ε_K , can be used to fit the meson charge ratios K^+/K^- and π^+/π^- . In subsequent sections we discuss how this model compares to several full Monte Carlo calculations of the muon charge ratio, and how a possible difference in the spectral index of the primary cosmic Hydrogen and Helium flux might affect the interpretations.

2 Previous Measurements of the Atmospheric Muon Charge Ratio

Numerous measurements of the atmospheric muon charge ratio data have been made at the earth's surface, from MeV to multi-TeV energies. The Hebbeker and Timmermans 2002 compilation article [5] provides many references to these data. We have compiled selected data in Figure 1. We are focused on the muon surface energy range above 10 GeV. For the energy range below 100 GeV/c, we have only include the newer, precise L3+C data sets [6]. As discussed below, we have compiled ratio data above 100 GeV. The ratio is well known to be about 1.2 - 1.4 above a few GeV. High precision data are critical to understanding the behavior of the ratio with energy.

The CosmoALEPH experiment [7] at the LEP facility at CERN published 8 data points in the 80-1600 GeV surface energy range. The cosmic ray portion of the experiment used only the hadron calorimeter and the time projection chambers. The detector was located at a depth of 320 meters of water equivalent (mwe). Apparently the magnetic field was not reversed during data collection. The zenith range of 0-10 deg was studied, but the publication did not give the ratio as a function of zenith angle.

Rastin [8] published data in the muon surface energy range from 6 - 1288 GeV using the Nottingham solid iron magnetic spectrometer. Data was collected with both forward and reversed magnetic fields.

Baxendale et al. [9] published extensive muon flux data in the momentum range 7-500 GeV/c using the Durham spectrometer, consisting of magnetized iron blocks, flash tubes, and scintillation counters. The magnetic field in the three magnetized blocks was reversed regularly to reduce bias effects. Their highest energy charge ratio data point is at 358 GeV/c. The publication did not give the ratio as a function of zenith angle.

The L3+C experiment has published extensive muon charge ratio data at many zenith angles at momentum up to 380 GeV/c. The high accuracy and extensive angular range of this data is ideal for comparison with our parameterization of the muon charge ratio. We have used this data extensively in Section 5 of this paper.

Matsuno et al. [10] published data in the momentum range from 70 GeV/c to 20 TeV/c using the MUTRON magnetic cosmic ray spectrometer. It consists of a solid iron magnetic spectrometer, a calorimeter of 16 proportional chambers, and 48 spark chambers. Data was collected with both forward and reversed magnetic fields. The data was collected only at large zenith angles 86-89 degrees. While the data sample is large, there are only a few hundred events above 2 TeV/c; these higher energy data will not be included in this analysis. Only the energy dependence of the charge ratio is used by us.

Experiments with data points with relatively large error bars have not been included in this analysis since they are not useful for the precise studies in this paper. Experiments with charge ratio data values above 100 GeV, with all data points having quoted errors greater than ± 0.1 , have not been included in our analysis; thus Hayman




Figure 1: Compilation of muon charge ratio from experiments with at least one data point with an uncertainty less than 0.10. The Hebbeker line is the fit of Hebbeker et al. to their compilation of low energy, near vertical cosmic muon data.

and Wolfendale [11], Appleton et al.[12], Nandi et al.[13], and Kremer et al.[14] are not included. Specifically, Burnett et al. [15] was not included in our study due to their absence of any published estimates of systematic error on their ratio values.

Using the above-discussed data, and the MINOS near detector and far detector data, Figure 1 displays the muon charge ratio data from 0.1 - 10 TeV muon surface energy $E_{\mu,0}$. The ratio is near 1.25 at low energies, but the MINOS data above 1 TeV is near 1.4. The MINOS Far Detector data clearly shows the rise of the ratio with energy. The rise in the ratio above 100 GeV has been attributed to the increasing contribution from Kaons[2]. The data in Figure 1 could be parameterized with a linear function of $E_{\mu,0}$ or a $\log(E)$ function. But there is no physics reason to expect a linear dependency with E , or even a $\log(E)$ dependency. We have chosen to use a different parameterization that encompasses more of the kinematics and the physics of the muon production processes. The detailed physics of this rise with energy is the subject of the analysis in Section 5.

3 Issues in measuring the Charge Ratio

In this section, we review the method used for determining the muon charge in a magnetic detector, and consider some of the systematic errors which affect such a measurement. We describe the concept of Maximum Detectable Momentum, and consider two distinct types of systematic errors: bias and randomization.

Most measurements of the atmospheric muon charge ratio involve the use of a magnet to deflect the trajectory of a charged particle. The resulting curvature is used to measure both the charge sign and momentum of the muon. Depending on the size, granularity and strength of the magnetic field, every detector has a Maximum Detectable Momentum (MDM). We define the MDM as that momentum for which a nearly straight real track will have a measured curvature (determined from a fit to points along the track) which is one standard deviation from zero. For a uniform magnetic field, the MDM is simply the reciprocal of the error (s.d.) of the curvature measurement, when the curvature is expressed in $(\text{GeV}/c)^{-1}$.

The MDM can be calculated for each event in each detector. As a simplified by instructive example, we consider a simplified toroid detector which is a long right circular cylinder of radius R , with uniform density, and the B field is azimuthal and constant. The long z -axis is perpendicular to the zenith direction. We do not consider any tracks which enter the front or leave the rear of a finite length cylinder, i.e. all muons enter and exit the side. All such μ tracks will cross near the center of the detector and be roughly S shaped, with curvature one way for the first half of the track, and the other way for the second half. The length of the corresponding half-track is

$$L = \sqrt{R^2 - b^2} / \sin(\theta) \quad (3)$$

where b is the impact parameter or distance of closest approach to the center of the toroid, and θ_z is the angle of the track with the z axis (not the zenith angle). For this discussion, b is near zero and θ is near 90 degrees. To calculate the precision of momentum measurement, we need the component of the magnetic field B which is perpendicular to the track direction. Approximating at the midpoint of the half-track, assuming that theta is not small, we find

$$B_{PERP} = B \times \sqrt{[1 - \sin^2(\theta) \times \cos^2(\arctan(\sqrt{(R^2 - b^2)/2b}))]} \quad (4)$$

This is used in Reference [16] to calculate the MDM for MINOS, which varies dramatically between 470 GeV to as low as 15 GeV for large impact parameters and large θ_z , where many muon tracks occur.

The above discussion deals with the ability to measure charge in an ideal detector. In a real detector, there are a number of reasons that we might assign an incorrect charge sign to a track. We separate these systematic errors into two kinds which we call randomization and bias. One kind of error involves biases towards one sign or the other. Other errors will cause tracks to be randomly assigned as positive or negative. These randomized events will have an apparent charge ratio of unity. Since the true charge ratio is greater than one, any such errors will lower the measured charge ratio. We next describe two instances of each kind of error.

The first kind of bias is associated with the magnetic geometry of the detector, which we call acceptance effects. Ignoring the very small magnetic field in the air outside the detector, the charge ratio of tracks which enter the detector is not affected by the magnetic field and/or the geometry of the detector. Instead, there are effects which have to do with cuts that are made to require a good track, and the efficiency of those cuts. From a given direction, tracks of one kind (the other) will be bent such that the track length in the detector is longer (shorter). In order to require a minimum number of hits to make a good charge determination, a minimum length cut may be made. Thus tracks in some directions may be biased towards one sign or the other. In the

MINOS geometry, such directions occur as peaks or dips in the azimuthal distributions of the charge ratio. They tend to be symmetric with respect to the detector, but since a varying overburden can cause more tracks from some directions than others, they do not necessarily cancel.

A second kind of bias comes from any misalignment of detector planes which causes a curvature in the coordinate system. A result of any such misalignment would be that straight tracks appear to have curvature. With the MINOS geometry, it was found that a systematic misalignment of even a fraction of a mm (as measured by the track's sagitta) would have a noticeable effect on the charge ratio. A curvature cut does not remove such a bias.

Bias effects as described above are correctable in principle, with an accurate Monte Carlo. Acceptance issues can be corrected if the model of the magnetic field is accurate, and if the overburden is correctly modeled so that the expected fluxes as a function of zenith and azimuthal angle are known. Misalignments can be modeled in principle, but must be done with a correct three-dimensional model of the detector, and must be done with an accuracy that is difficult to achieve in practice.

An easier way to deal with the above bias effects is to reverse the magnetic field. If the reversal of the field is done with sufficient accuracy, then all forms of bias described above cancel when one uses the geometric mean of the charge ratio for forward and reversed field running

$$r = [r_{forward} \times r_{reversed}]^{1/2} \quad (5)$$

This cancellation is apparent when one considers the ratio of the positive muons in forward(reverse) running to negative muons in reverse(forward) running. The equality of these two values is a powerful consistency check.

By contrast, effects which tend to randomize the charge ratio do not typically cancel in this manner and must be dealt with separately. An obvious type of randomization occurs for high momentum straight tracks above the MDM. If a track is straight within the resolution of the detector, then we will be unable to accurately determine the charge sign. For a straight enough track, a program will measure curvature either way an equal fraction of the time. The fit that calculates the curvature can also calculate the error on the curvature. This form of randomization can be minimized by cutting on the fractional curvature error $\sigma(1/p)/(1/p)$ as is shown in Figure 5 of Reference [1]. However the fraction of mismeasured tracks may not follow a Gaussian distribution for $\sigma(1/p)/(1/p)$ and Monte Carlo estimates of the misidentification must be analyzed with care.

A second form of randomization, which was observed in MINOS data, was generally associated with extra hits. These might be due to noise, crosstalk or demultiplexing issues, or also delta rays or other particles associated with the muon. For beam experiments, the magnetic field is perpendicular to the beam direction, to maximize bending power from the magnet. For cosmic rays, muons are observed at all angles. As described above, muon tracks will be bent much more in a toroidal magnet if they impinge on the detector with a low impact parameter. For those cosmic ray muons which are traveling mostly parallel to the magnetic field, an interesting situation can arise if the apparent bending due to multiple scattering is comparable to the bending from the magnetic field. If that happens, a reconstruction program which takes into account multiple scattering can get a good fit for a straight track for any momentum. The inclusion of a hit or hits which do not belong on the track can then lead to a class of muon tracks which are really at high momentum, but get reconstructed at low

momentum with a random charge sign depending on which side of the track the extra hits appeared.

We can quantify this result as follows: Projected onto a plane, the RMS multiple scattering angle can be written

$$\langle \theta_{MCS} \rangle = [0.0136/p(\text{GeV}/c)] * \sqrt{(L(\text{meter})/X)} \quad (6)$$

where X is the radiation length (0.0383 m in MINOS). The magnetic bending for this same length of track is given by

$$\theta_{Bend} = L(\text{meter}) * B_{PERP}(\text{Tesla})/[3.336 \times p(\text{GeV}/c)] \quad (7)$$

Note that in the ratio, p cancels. For typical muon tracks in MINOS, the ratio rises from 0.14 at normal incidence to near unity at large impact parameter. Thus the type of randomization described in the previous paragraph can be preferentially removed by throwing away events with large values of $\langle \theta_{MCS} \rangle / \theta_{Bend}$.

The cancellation of bias effects can be checked by studying distributions which have bias effects in them before and after the forward and reverse field data are combined. We have no such consistency check for randomization effects. Thus, the ability to cancel all bias effects using the combined forward and reverse field data is stronger than the ability to cancel all randomization effects. This might lead to a tendency to report a charge ratio which is systematically smaller than the true value. We also note that not every experiment listed in Section 2 has taken data with reversed magnetic field.

4 Issues with Interpreting Data Obtained Underground

4.1 The relevance of energy loss differences

The charge ratio that MINOS and other underground experiments have measured is a ratio of μ^+ to μ^- at the detector. For comparison with production models, the relevant quantity is the ratio in the upper atmosphere. If the energy loss of positive and negative muons were identical, as they go through the earth, these two ratios would be the same. The leading energy loss processes of μ^+ and μ^- are the same of course, but there are corrections of the order of the fine structure constant α . We evaluate these small differences, first for ionization energy loss, and then for radiative processes. The latter difference is found to be negligible. The small difference in ionization energy loss is used to calculate the difference in range and the correction to the charge asymmetry as a function of slant depth in Section 4.4.

The statistical energy loss of muons, traversing an amount X of matter in g/cm^2 , with energies far above the Bethe-Bloch minimum is usually parameterized as

$$-\frac{dE_\mu}{dX} = a(E_\mu) + \sum_{n=1}^3 b_n(E_\mu) \cdot E_\mu, \quad (8)$$

where a is the collisional term (i.e. ionization, mostly due to delta-ray production) and b in the second term accounts for the three radiative muon energy loss processes: 1. Bremsstrahlung and 2. pair production, and 3. photo-nuclear interactions. In Table

[17][18] these energy loss parameters are listed for standard rock. The critical energy where ionization losses equal radiative losses in standard rock is approximately 0.6 TeV. The average muon surface energy for a muon which reaches the depth of MINOS is greater than 1 TeV, so the b term and its energy dependence are important in calculating the energy loss. We have investigated differences in the a and b terms for μ^+ and μ^- .

E_μ [GeV]	a_{ion} [MeV cm ² /g]	b_{brems}	b_{pair}	b_{DIS}	Σb
		[10 ⁻⁶ cm ² /g]			
10	2.17	0.70	0.70	0.50	1.90
10 ²	2.44	1.10	1.53	0.41	3.04
10 ³	2.68	1.44	2.07	0.41	3.92
10 ⁴	2.93	1.62	2.27	0.46	4.35

Table 1: Average muon energy loss parameters calculated for standard rock [17][18]

4.2 Difference in Ionization dE/dx for μ^+ and μ^-

At low energies, around the Bethe-Bloch maximum, the difference in ionization energy loss is known as the Barkas effect[19], and there have been efforts to both measure and calculate those differences[20]. Calculations show that negative particles lose energy at a slower rate, with the difference dropping from tens of percent at MeV energies to about 0.3% in the GeV range. Such differences were experimentally verified both at MeV energies[21][22][23] and in the GeV range [24]. At higher energies, this difference in ionization energy loss has usually been neglected, and we are not aware of any measurements. As described in Reference [25], the usual ionization energy loss term for muons (of either sign) depends on z^2 , and the difference between μ^+ and μ^- arises from a small additional z^3 correction term. This correction term in dE/dx is:

$$\left(\frac{dE}{dX}\right)_{ion}^{corr} = \frac{\pi\alpha z^3 0.307Z}{2\beta A} \quad [MeV cm^2 g^{-1}] \quad (9)$$

where α is the fine structure constant, z is the charge, β is the relativistic velocity, and Z and A are the nuclear properties of the material through which the muon is passing. The absolute value of the difference in ionization energy loss between positive and negative muons in standard rock[18] is plotted in Figure 2. It is fairly constant above 10 GeV, at a value corresponding to approximately 0.15% of the mean energy loss in the ionization dominated energy regime (*c.f.* Table 1).

4.3 Calculated Difference in Bremsstrahlung dE/dX for μ^+ and μ^-

Above an energy near 0.6 TeV in standard rock, radiative energy loss becomes comparable to ionization energy loss, and continues to grow at higher muon energies. From Reference [26], the fractional difference in Bremsstrahlung energy loss between positive and negative muons is

$$\frac{[\frac{dE}{dX}]_{brems}^{\mu^+} - [\frac{dE}{dX}]_{brems}^{\mu^-}}{[\frac{dE}{dX}]_{brems}} = \frac{8Z\alpha}{\gamma} \quad (10)$$




Figure 2: Calculated difference in ionization energy loss between positive and negative muons in standard rock (average nuclear properties: $\bar{Z} = 11$, $\bar{A} = 22$ [18]).




Figure 3: Calculated fractional difference in Bremsstrahlung energy loss between positive and negative muons.

where γ is the Lorentz factor of the muon. Again, the μ^+ has a slightly higher energy loss. This fractional difference *decreases* with energy and is already negligible where radiative energy losses become important. This fractional difference is plotted in Figure 3 for muons in standard rock. Presumably, the same fractional difference can also be assigned for pair-production, as the underlying process is a two-photon exchange between the muon and the constituents of the nucleus, and thus the cross sections for μ^+ and μ^- should scale in the same way as for Bremsstrahlung.

4.4 Range Underground and muon charge Asymmetry

Taking the vertical muon intensity from the Gaisser parameterization of the muon flux at the surface (see Equation 2) and propagating this energy spectrum underground according to statistical ionization and radiative energy losses, it is possible to precisely calculate the underground muon intensity. This procedure is described in detail in [27] for overburdens of standard and Soudan rock (MINOS). First, the average muon




Figure 4: Calculated ratio of positive to negative vertical muon intensities in Soudan rock as a function of slant depth for equal surface intensities. The upper curve is for all muons, the lower curve is for muons with a remnant momentum of less than 250 GeV/c (\approx the maximum detectable momentum in the MINOS far detector).

range underground, for each value of surface energy, is computed. For this, the energy dependent a and Σb values were parameterized for standard and Soudan rock as in [27]. The additional ionization loss according to Equation 9 was added for μ^+ to the value of the function for a (subtracted for μ^-). Radiative losses (90 % of Σb) are scaled with the energy dependent fractional difference $R = 8Z\alpha/\gamma$ according to Equation 10 as $\Sigma b^\pm = 0.9 \cdot \Sigma b \cdot (1 \pm R/2)$ for μ^+ and μ^- , respectively. The contribution from photo-nuclear production (DIS), is not scaled. For each initial value of muon energy, the slant depth in meter-water-equivalent where the muons of different charge range out is determined.

We compute the underground intensities of positive and negative muons as a function of slant depth for a given rock composition. This ratio of the μ^+ and μ^- intensity curves is shown in Figure 4 for Soudan rock. The upper curve corresponds to the fractional difference in integral intensities of μ^+ and μ^- at a given slant depth. For slant depth values above about 1000 mwe the underground ratio $N(\mu^+)/N(\mu^-)$ is lowered by roughly 0.4 %. The detected intensity corresponds to the charge ratio of the muons at depth below the MDM for MINOS. For increasing slant depth values the measured underground ratio $N(\mu^+)/N(\mu^-)$ is further reduced and saturates at about 0.6 % below its surface value for slant depths larger than roughly 5000 mwe.

The small fractional difference in energy loss for μ^+ and μ^- of the order of 0.15 % predicted by theoretical calculations at high energies, gets amplified by a factor of about 3.7, due to the approximate $E^{-3.7}$ dependence of the differential muon spectrum. The impact of the rock composition is almost negligible, as the induced muon charge asymmetry under Soudan rock lowers the surface value of the ratio by an additional amount less than 0.02 % compared to standard rock. Figure 4 can be used to correct the underground measured muon charge ratio to its surface value. For MINOS, this corresponds to a correction which increases the measured value by 0.6%.

5 Model of Pion and Kaon Contributions to the Charge Ratio

We have investigated a generalization of Gaisser's Equation 2 to study separately the positive and negative muon intensities. Using the positive fraction parameters f_π and f_K , the energy dependency of the positive and negative muons is given by

$$\frac{dN_{\mu^+}}{dE_\mu} = \frac{0.14E_\mu^{-2.7}}{\text{cm}^2 \text{ s sr GeV}} \times \left\{ \frac{f_\pi}{1 + \frac{1.1E_\mu \cos \theta}{115 \text{ GeV}}} + \frac{\eta \times f_K}{1 + \frac{1.1E_\mu \cos \theta}{850 \text{ GeV}}} \right\} \quad (11)$$

$$\frac{dN_{\mu^-}}{dE_\mu} = \frac{0.14E_\mu^{-2.7}}{\text{cm}^2 \text{ s sr GeV}} \times \left\{ \frac{1 - f_\pi}{1 + \frac{1.1E_\mu \cos \theta}{115 \text{ GeV}}} + \frac{\eta \times (1 - f_K)}{1 + \frac{1.1E_\mu \cos \theta}{850 \text{ GeV}}} \right\} \quad (12)$$

where the values for ϵ_π and ϵ_K have been replaced by their numerical values. We can use equations 11 and 12 to calculate the surface muon charge ratio:

$$r_\mu = \frac{\left\{ \frac{f_\pi}{1 + 1.1E_\mu \cos \theta / 115 \text{ GeV}} + \frac{\eta \times f_K}{1 + 1.1E_\mu \cos \theta / 850 \text{ GeV}} \right\}}{\left\{ \frac{1 - f_\pi}{1 + 1.1E_\mu \cos \theta / 115 \text{ GeV}} + \frac{\eta \times (1 - f_K)}{1 + 1.1E_\mu \cos \theta / 850 \text{ GeV}} \right\}} \quad (13)$$

The charge ratio of muons from pion decay is $r_\pi = f_\pi / (1 - f_\pi)$ and from Kaon decay is $r_K = f_K / (1 - f_K)$. We will refer to the implications of Equation 13 with energy independent parameters as the pika model. There are several interesting features of this model:

1. The relative intensity of cosmic ray pions and Kaons that contribute to muon production can be extracted from surface and underground muon charge ratio experiments.
2. The muon charge ratio does not depend upon the muon energy and the zenith angle separately, but on the product $E_\mu^{sur\ face} \cos \theta_z$. This product of terms controls the relative portions of interaction and decay for both pions and Kaons. At a fixed value of $E_\mu^{sur\ face} \cos \theta_z$, the intensity ratio of muons from pions and Kaons is constant.
3. There are contributions to the muon charge ratio from both pions and Kaons from 1 GeV to 10 TeV. The contribution to the ratio from Kaons does not vanish at GeV energies - it is just smaller than at TeV energies. This effect can be seen in Figure 5 which displays the contribution to the muon flux from each of the four mesons.
4. The pika model postulates an energy independent π^+ / π^- ratio related to f_π , and energy independent K^+ / K^- ratio related to f_K , and an energy independent π / K ratio embodied in the Gaisser constant 0.054.
5. Contributions from charm particle production have been ignored because their effect on the ratio below 10 TeV is expected to be negligible.

The energy independence of η , f_π and f_K is consistent with Feynman scaling. Here we explore features of our simple model, and in the next section we shall compare the qualitative features of our model with full simulations.




Figure 5: Contributions to the muon flux from the four mesons using the pika model

In our parameterization, the muon surface charge ratio depends only on the relative number of pions and Kaons that yield muons which are detected on the surface (or underground). The relative contribution to the muon intensity from different mesons is affected by the relative probability of hadron interaction and decay, which vary with both meson energy and zenith angle. The intensity varies with energy because of the hadron lifetimes, and it varies with angle because of the changing interaction probability in the atmosphere. However, inspection of equation shows that at any fixed value of $E_{\mu}^{surface} \cos \theta_z$, the ratio of muons from pion and Kaon decay is also fixed, even if E or θ is varied.

Our parameterization also yields additional insight into the sensitivity of various underground experiments which measure the muon charge ratio. Underground experiments can effectively observe a much smaller range of $E_{\mu}^{surface} \cos \theta_z$ than a range of E. To see this, consider a simple approximation with five assumptions:

1. The parameter a in Equation 8 is constant
2. The parameter b in Equation 8 is constant and set to zero.
3. The earth/rock density above the underground detector is constant
4. The surface above the detector is flat
5. The MDM in the underground detector is negligible compared to the energy loss from the surface to the detector.

The first four assumptions are equivalent to stating that the muon energy loss in the earth is proportional to the overburden which only depends upon the muon zenith angle. In that case the energy loss through the overburden is $E_{loss} = E_{min}/\cos(\theta)$, where E_{min} is the minimum energy loss for a vertical cosmic muon. The fifth assumption implies that the underground energy of the muons used for the charge ratio measurement is much smaller than their surface momentum.

With these five assumptions, the muon intensity distribution in $E_\mu^{surface} \cos \theta_z$ would be measured to be a delta function at a value of E_{min} . While at large zenith angles the surface energy increases due to the $1/\cos(\theta)$ dependence of the slant depth, the combined quantity $E_\mu^{surface} \cos \theta_z$ remains constant. It is illuminating to compare this predicted behavior to the actual distribution in the MINOS detectors when no such assumptions are made. These are shown in Figures 7 and 6. Both of these MINOS $E_\mu^{surface} \cos \theta_z$ distributions are considerably narrower than the corresponding $E_{surface}$ distributions. The $b(E)$ radiative term yields the largest contribution to the width of the measured $E_\mu^{surface} \cos \theta_z$ distribution for the MINOS far detector. In the near detector distribution, the largest contribution is the larger ratio of maximum detectable momentum to energy loss in the overburden.




Figure 6: Distribution of E and $E_\mu^{surface} \cos \theta$ for MINOS data muons in the Far Detector, after cuts (GeV).




Figure 7: Preliminary distribution of E and $E_\mu^{surface} \cos \theta$ for MINOS data muons in the Near Detector, after cuts (GeV). Note that the $E_\mu^{surface} \cos \theta$ distribution is so narrow that it goes far off scale, but there are the same events in each distribution.

We have used Equation 13 and the measured muon charge ratio to study r_π and r_K . We have done chi-squared fits in $E_\mu^{surface} \cos \theta_z$ to the MINOS near detector and far detector data, and to the L3+C data. The fit yields $f_\pi = 0.555 + -0.002$ and $f_K = 0.667 + -0.007$. These values lead to a muon charge ratio from pion decay of $r_\pi = 1.25$ and a muon charge ratio from charged Kaon decay of $r_K = 2.0$. The errors in the two parameters are highly correlated. The fits are shown in Figure 15 along with the MINOS and L3+C data. The agreement between the parametrization and the data is excellent.

Using the f values from this fit, we plot Equation 13 for an extended range of muon energy in Figure 8. The critical energies discussed above for pion and Kaons are indicated by the arrows. The low energy asymptotic ratio is $(f_\pi + \eta f_K) / [1 - f_\pi + \eta(1 - f_K)] = 1.26$. The high energy ratio is $(\varepsilon_\pi f_\pi + \varepsilon_K \eta f_K) / [\varepsilon_\pi(1 - f_\pi) + \varepsilon_K \eta(1 - f_K)] = 1.45$. The low and high energy asymptotic values each consist of a different ratio of muons from π decay and from K decay.

The fit was repeated using just the smaller data sample of the MINOS near detector [28] and far detector, the fit was repeated. We obtain $f_\pi = 0.549 + -0.012$ and $f_K = 0.702 + -0.049$, which imply $r_\pi = 1.22$ and $r_K = 2.36$. The errors on the two parameters are highly correlated.




Figure 8: The pika model’s muon charge ratio for an extended range of energy. The critical energies for pion and Kaon decay vs interaction are shown by the arrows.

The parameterization we have used seems sufficient to represent the published data sets. A parametrization linear in $E_{\mu}^{surface}$ or $\log E_{\mu}^{surface}$ is less well motivated.

Our fits to r_{π} gives values near expectations. Our fit to $r_K = K^+/K^-$ in atmospheric showers yield values just above 2.0. It is clearly difficult to directly measure the Kaon charge ratio. Equation 13 provides a well defined parametrization for future studies of this subject.

6 Comparison with Full Simulations of Atmospheric Muons

The three full simulations of CORT [29], Lipari[30], and Honda [31] have published the charge ratio as a function of E for various ranges of $\cos(\theta)$. These model results versus $E_{\mu}^{surface}$ are displayed in Figures 9, 10, and 11. Results versus $E_{\mu}^{surface} \cos \theta$ are shown in Figures 12, 13, and 14.

Independent of any data, we investigated (1) the dependency of these ratio simulations on the $E_{\mu}^{surface} \cos \theta_z$ variable, and (2) their consistency with the pika model of the charge ratio. Our observation will be that $E_{\mu}^{surface} \cos \theta$ is a more useful variable for ratio analysis than just the muon energy $E_{\mu}^{surface}$. We also find that simulations are using a better value of r_{π} than r_K .

Figure 12 displays Honda’s muon charge ratio simulations as a function of $E_{\mu}^{surface} \cos \theta_z$ for seven ranges of $\cos(\theta)$ from 0.3 to 1.0. While there are certainly variations in the simulations as the angle changes, most of the variation seen in Figure 12 is suppressed up to ~ 4 TeV. The dip in the charge ratio near ~ 70 GeV has probably the strongest angular dependency. Above ~ 200 GeV, the charge ratio is nearly independent of




Figure 9: Honda calculations of the muon charge ratio at 5 zenith angles




Figure 10: CORT calculations of the muon charge ratio at 4 zenith angles




Figure 11: Lipari calculations of the muon charge ratio at 5 zenith angles

$\cos(\theta)$, as predicted by the pika model.

Figure 13 displays the CORT theory simulations for four values of $\cos(\theta)$. In the $E_{\mu}^{surface} \cos \theta_z$ range from 5 GeV to 600 GeV, the simulations show little dependency on $\cos(\theta)$. Above a TeV, the ratio does increase with $\cos(\theta)$. However, ~ 1 TeV is near the high energy limit of this simulation. Above 600 GeV, the charge ratio increases with increasing values of $\cos(\theta)$.

Figure 14 displays the Lipari simulations for five values of $\cos(\theta)$ from 0.2 to 1.0. It is apparent that a significant portion of the predicted charge ratio variation with $\cos(\theta)$ is suppressed when the ratio is calculated as a function of $E_{\mu}^{surface} \cos \theta_z$.

The observed dependence of the above three simulations on $E_{\mu}^{surface} \cos \theta_z$ provides some justification for analyzing the charge ratio data using the pika formula. We would encourage the authors of the ratio simulations to understand the source of the remaining variations in their simulations as a function of $E_{\mu}^{surface} \cos \theta_z$ at values greater than ~ 1 TeV.

Next we directly compare the pika model to the three simulations discussed above. Two parameter chi-squared fits were performed to the simulations assuming equal uncertainties on each point within a simulation. The best fit for each simulation is shown as a smooth line in the corresponding figure. For the CORT model, the pika model fits well the simulation from 20 GeV to 600 GeV at all angles. The fit yields $f_{\pi} = 0.557$ and $f_K = 0.705$. For the Honda simulation, the pika model can not reproduce the simulation's dip at ~ 100 GeV, although it does describe the simulations energy dependency from 5 GeV to 5 TeV. For the Lipari simulation, the pika model reproduces well the simulation from 3 GeV to 10 TeV. The agreement at low energy is expected.

Next we have compared the L3+C and MINOS charge ratio data sets to the three same simulations. To summarize the three simulations, we have used the fits of the




Figure 12: Honda calculations of the muon charge ratio vs $E_{\mu}^{surface} \cos \theta_z$ at 4 zenith angles. The curve is a fit of these simulation results to the pika model

three simulations to the pika model to compare with data. As seen in Figure 16, none of the models (as parameterized by the pika formula) correctly predicts the data's dependency with $E_{\mu}^{surface} \cos \theta_z$. The simplest explanation is that the simulations have not used the correct value for r_K . This is calculated in Table 3 and discussed further in Section 8.

7 Effect of Helium on the Charge Ratio

Another process that could affect the energy dependence of r_{μ} would be a different spectral index for heavier cosmic ray primaries than for Hydrogen. This would introduce an energy dependence to the incoming proton to neutron ratio. It is not well established that the energy dependency of the heavy primaries is different than that for hydrogen in the TeV energy range. For this study, we will use Gaisser and Honda's [32] parameterizations of the primary flux as a function of the primary component energy E_k , given as

$$\phi(E_k) = K(E_k + b \times \exp[-c\sqrt{(E_k)}])^{-\alpha} \quad (14)$$

The parameters in the above equation are given in the Table 2 below. Note that this reference has a spectral index decreasing slightly with increasing primary mass.

Figure 17 displays the flux as a function of the kinetic energy per nucleon. Based on Monte Carlo calculations using CORSICA, the fraction of the primary proton energy that is transferred to the muon is approximately 9%. The corresponding fraction of the primary helium energy, per nucleon, that is transferred to the muon is approximately 13%. Without using a simulation we will assume the fraction of the energy transfer for




Figure 13: CORT calculations of the muon charge ratio vs $E_{\mu}^{surface} \cos \theta_z$ at 4 zenith angles. The curve is a fit of these simulation results to the pika model




Figure 14: Lipari calculations of the muon charge ratio vs $E_{\mu}^{surface} \cos \theta_z$ at 5 zenith angles. The curve is a fit of these simulation results to the pika model




Figure 15: pika model compared to L3+C and MINOS data sets




Figure 16: Fits to the three simulation models of ratio vs $E_{\mu}^{surface} \cos \theta_z$ compared to L3+C and MINOS data sets

Parameter	α	K	b	c
Hydrogen	2.74 ± 0.01	14900 ± 600	2.15	0.21
He (A=4)	2.64 ± 0.01	600 ± 30	1.25	0.14
CNO (A=14)	2.60 ± 0.07	33.2 ± 5	0.97	0.01

Table 2: Primary flux parameters used in the text.




Figure 17: Fractional contribution of various nuclei to the nucleon spectrum

primary carbon, nitrogen, and oxygen to also be 13%. When the energy profiles of the protons and Helium from Monte Carlo are used instead of these average values, the impact on the charge ratio is observed to be not large. For convenience and clarity, we will use the average values in the analysis in this section of the paper.

We will model the change in the muon charge ratio due to just pion decay when helium and CNO are added to the primary proton flux. In a simple quark model the leading proton and neutron particle in the forward direction will give rise to a charge ratio of 1.00 for symmetric nuclei $A-Z=Z$. Setting the muon charge ratio equal to 1.25 at 1 GeV and fixing the normalization of the hydrogen flux to be 1.32 (in order to get back the charge ratio of 1.25 at 1 GeV), one has for the muon charge ratio

$$R^\pi(E_\mu) = \frac{1.32 \times \phi_p(E_\mu/0.09) + 1 \times 4 \times \phi_{He}(E_\mu/0.13) + 1 \times 14 \times \phi_{CNO}(E_\mu/0.13)}{\phi_p(E_\mu/0.09) + 4 \times \phi_{He}(E_\mu/0.13) + 14 \times \phi_{CNO}(E_\mu/0.13)} \quad (15)$$

Figure 18 displays the corresponding fractional change in the charge ratio as a function of the muon surface energy. Note that the effect is more than a 3 percent reduction at 1 TeV. This dependency can be fit to a polynomial in $\log(E)$, giving the expression




Figure 18: Fractional change of R

$$R_{rel}^{par}(E_\mu)_\pi = 1 - 0.00575 \times \log_{10}(E_\mu) - 0.00175 \times \log_{10}^2(E_\mu) \quad (16)$$

Next we include the contribution from Kaon production by the primaries. Again assuming a quark model for the leading Kaon in the forward direction from incident protons and neutrons, one has

$$p(uud) + air \rightarrow K^+(u\bar{s}) + \dots \quad (17)$$

and

$$n(duu) + air \rightarrow K^+(u\bar{s}) + \dots \quad (18)$$

This leads to a charge ratio of

$$1 + 3/4\epsilon_{K^+}^p \quad (19)$$

for symmetric nuclei with $A-Z=Z$. We further assume that the muon charge ratio at 1 TeV is 1.374 (near the MINOS value). This then yields

$$R^K(E_\mu) = \frac{2.7 \times \phi_p(E_\mu/0.09) + 2.3 \times 4 \times \phi_{He}(E_\mu/0.13) + 2.3 \times 14 \times \phi_{CNO}(E_\mu/0.13)}{\phi_p(E_\mu/0.09) + 4 \times \phi_{He}(E_\mu/0.13) + 14 \times \phi_{CNO}(E_\mu/0.13)} \quad (20)$$

The effect on the muon charge ratio due to the heavy ions requires that the pion and Kaon fractions be modified as follows:

$$f_\pi^*(E_\mu) = 1/(1 + 1/r_\pi \times R(E_\mu)) \quad (21)$$

$$f_K^*(E_\mu) = 1/(1 + 1/r_K \times R(E_\mu)) \quad (22)$$

Including these contributions from heavy primaries does have an impact on the calculation of parameters of the pika model. The charge ratio no longer depends just




Figure 19: Fit to the meson charge ratios modifying the pika model with the energy dependence of helium production.

upon $E_\mu \cos \theta$. While this effect is small it is certainly present and can be accounted for in modeling. Figure 19 shows the effect of the heavy primaries on the two parameters of the pika model. The fit to the data is almost indistinguishable from the previous fit. Note that an increasing heavy fraction at high energy will decrease the charge ratio, so to fit to the high MINOS points, a larger value of r_K is required. It is clear that simulations of the muon charge ratio need to include the possible different energy dependence of heavy primaries.

We repeat that the choice of spectral index in Table 2 is to illustrate the size of a possible effect. If it turns out that the spectral index is independent of chemical composition, the effect described in this section will not exist.

8 Summary and Discussion

We have reviewed several factors which affect the measurement and interpretation of the muon charge ratio r_μ deep underground. While many quantities in cosmic ray physics are difficult to measure precisely, r_μ has been measured in MINOS with a statistical accuracy better than 0.3%. In order to take advantage of this high precision, an experiment has to control systematic errors to a comparable level, which can be challenging. We pointed out two kinds of systematic error, those which might bias the measurement of r_μ , and those which randomize, and hence lower the measured value from the true value. Bias errors can be canceled to high precision by using data with both magnetic field polarities and Equation 5, and the success of this cancellation can be checked for consistency. Randomization errors, on the other hand, may require severe cuts that affect the statistical precision. Variables which may be related to

randomization effects can be identified and used to cut out data which have random background. However there is no independent way to determine if all such effects have been eliminated. The possibility exists that the true values of the charge ratio are higher than those that have been reported.

The MDM is an important parameter for an experiment with a magnet. It affects both the calculation of the systematic error and the energy resolution, and also directly limits the range of detected muon energies useful for r_μ and $E_\mu^{surface} \cos \theta$.

We have considered the change in the charge ratio from the fact that μ^+ lose slightly more energy than μ^- while crossing the overburden of an underground detector, both for ionization and catastrophic energy loss. For catastrophic energy loss, the effect is negligible, even for the small errors in MINOS. For ionization, we have calculated that the 0.15% difference in energy loss leads to a 0.6% difference in measured r_μ at 2000 mwe. This effect was mentioned in Reference [1] but not used to correct the reported charge ratio there.

The central result of this paper is the development and application of Equation 13 and the consequent relationship of r_μ to r_π and r_K . We have used the equation to study its consistency with several simulations, and to make fits to data. In Table 3, we show the fits for our parameters to the simulations. We also show the fits to the data reported in References [1], [6] and [33].

		r_K	r_π
Simulations	Lipari	2.87	1.25
	CORT	2.39	1.26
	Honda	1.63	1.28
	Gaisser	3.2	1.4
	Agrawal	2.92	1.35
Data fits	MINOS and L3	2.03	1.22
	MINOS N+F	2.23 ± 0.26	1.232 ± 0.030
	global	2.28 ± 0.06	1.224 ± 0.003
	Helium 2.74 \rightarrow 2.64	2.8	n/a

Table 3: Fits to r_K and r_π for data and simulations

We have shown the importance of using $E_\mu^{surface} \cos \theta_z$ (instead of just energy) in the analysis of the atmospheric muon charge ratio for energies above 10 GeV. Understanding of the muon charge ratio would benefit if future experiments and theory simulations would provide results using $E_\mu^{surface} \cos \theta_z$. The rise in the charge ratio, which occurs between the π and K critical energies of 115 GeV and 850 GeV, can be used to determine the meson charge ratios K^+/K^- and π^+/π^- . This is a new important technique for obtaining information on these ratios.

We further showed the effect on the charge ratio assuming that there is a spectral index for cosmic ray Helium nuclei which is different than the spectral index for protons. Using an index of 2.74 (2.64) for H (He), we get a change in r_K from 2.3 to 2.8 compared to an analysis where the spectral indices were the same.

Both the data and the simulations seem to have consistent values for r_π . A general feature of our analysis is that the new MINOS data by themselves suggest a lower value of r_K than is used in many of the simulations. However, it is important to note that the systematic effects considered in this paper, which include errors from randomization,

corrections for differences in dE/dx for μ^+ and μ^- and a possible lower spectral index for Helium, would all tend to raise the fitted values of r_K .

Several effects which have not been explicitly considered here are expected to be small. These include the production of muons from charm and other heavy particles, components of the cosmic rays heavier than helium and possible differences in their spectra, and a variety of scaling violations which would have the effect of making f_π and f_K energy dependent. These effects need full simulations to evaluate fully, but such simulations need the right average values of these parameters to be useful. It is in that context that the analysis presented in this paper can be useful.

We note that to fully exploit the rise in the charge ratio observed by MINOS, there is a need for additional precise ratio data in the $E_\mu^{surface} \cos \theta_z$ ranges of 0.2-0.8 TeV, and above 3 TeV.

9 Acknowledgments

This work was supported by the U.S. Department of Energy and Benedictine University. We would like to thank V. Naumov for introducing us to the issues involved in the predicted rise of the charge ratio. We thank Tom Gaisser, Morihiro Honda, Paolo Lipari and Teresa Montaruli for discussions on the ratio simulations. We acknowledge the help and insight of Giles Barr, Thomas Fields, Jeff de Jong and Alec Habig. We are grateful to Geoff Bodwin and Stanley Wojcicki for their contributions in understanding issues involving energy loss. We thank Eric Beall, Gavril Giurgiu, Eric Grashorn, Stuart Mufson, and Brian Rebel for their many contributions. And the support of the entire MINOS collaboration has been invaluable.

References

- [1] P. Adamson et al., "Measurement of the Atmospheric Muon Charge Ratio at TeV energies with MINOS", Phys. Rev. D76 052003 (2007).
- [2] G. Fiorentini, V.A. Naumov and F.L. Billante, Phys. Lett. B **510**, 173 (2001). V. Naumov, private communication.
- [3] T. Gaisser, "Cosmic Rays and Particle Physics", Cambridge University Press, 1990.
- [4] Zatsepin and Kuzmin, Soviet Physics JETP volume 12, June 1961
- [5] T.Hebbeker and C. Timmermans, Astropart. Phys. 18 107 (2002).
- [6] P. Achard et al., Phys. Lett. B 598, 15 (2004).
- [7] D. Zimmermann, et al., Nuclear Inst. and Methods in Physics Research A, 141, 525 (2004).
- [8] B. C. Rastin, J. Phys. G 1629 10 (1984).
- [9] J. M. Baxendale, C. J. Hume, and M.G.Thompson, J. Phys G 1, 81 (1975).
- [10] S. Matsuno et al., Phys. Rev. D 29, 1 (1984).
- [11] P.J. Hayman et al., Nature 195 166 (1962).
- [12] I. C. Appleton, M.T. Hogue, and B.C. Rastin, Nucl. Phys. B 26 365 (1971).

- [13] B. C. Nandi, M.S. Sinha, Nucl. Phys. B 40 289 (1972).
- [14] J. Kremer et al., Phys. Rev. Lett. 83 4241 (1999).
- [15] T. H. Burnett et al., Phys. Rev. Lett. 30 937 (1973).
- [16] M. Goodman for MINOS, “Maximum Detectable Momentum in MINOS”, Proceedings of the 30th International Cosmic Ray Conference, Merida Mexico, 2007.
- [17] D.E.Groom et al., “Muon stopping-power and range tables,” Atomic Data and Nuclear Data Tables, 78, 183 (2001).
- [18] W. Yao et al., J. Phys. G: Nucl. Part. Phys. **33** 2006; standard rock is described in section 24.1 on page 247.
- [19] Letter from E. Fermi to W. Barkas (1953).
- [20] J. Jackson and R. McCarthy, Phys. Rev. **B 6** 4131 (1972).
- [21] F. Smith, W. Birnbaum and W. Barkas, Phys. Rev. **91**, 765 (1953).
- [22] W. Barkas, W. Birnbaum and F. Smith, Phys. Rev. **101**, 778 (1956).
- [23] M. Agnello et al., Phys. Rev. Lett. **74** 371 (1995).
- [24] A. Clark et al., Phys. Lett. **41b** 229 (1972).
- [25] J. D. Jackson, Phys. Rev. **D 59**, 017301 (1998).
- [26] J. D. Jackson, private communication to S. Wojcicki, Nov. 4, 2005.
- [27] J. Reichenbacher, “Calculating the Crouch Curve”, Proceedings of the 30th International Cosmic Ray Conference, Merida Mexico, 2007.
- [28] J. DeJong for MINOS, “Measurement of the Atmospheric Muon Charge Ratio with the MINOS Near Detector”, Proceedings of the 30th International Cosmic Ray Conference, Merida Mexico, 2007.
- [29] V. Naumov, private communication.
- [30] P. Lipari, Astroparticle Physics 1 195 (1993).
- [31] M. Honda, et al., Phys. Rev. D 52 4985 (1995) and T. Sanuki et al., 29th International Cosmic Ray Conference 9 139 (2005).
- [32] Gaisser and M. Honda, arXiv:hep-ph/0203272 v2 (30 March 2002).
- [33] P. Schreiner and M. Goodman for the MINOS collaboration, “Interpretation of the Atmospheric Muon Charge Ratio in MINOS”, Proceedings of the 30th International Cosmic Ray Conference, Merida Mexico, 2007.